


Caenorhabditis elegans PRMT-7 and PRMT-9 Are Evolutionarily Conserved Protein Arginine Methyltransferases with Distinct Substrate Specificities

Andrea Hadjikyriacou¹ and Steven G. Clarke^{1*}

Department of Chemistry and Biochemistry and the Molecular Biology Institute, University of California, Los Angeles, 607 Charles E. Young Drive East, Los Angeles, California 90095-1569, United States

S Supporting Information

ABSTRACT: *Caenorhabditis elegans* protein arginine methyltransferases PRMT-7 and PRMT-9 are two evolutionarily conserved enzymes, with distinct orthologs in plants, invertebrates, and vertebrates. Biochemical characterization of these two enzymes reveals that they share much in common with their mammalian orthologs. *C. elegans* PRMT-7 produces only monomethylarginine (MMA) and preferentially methylates R-X-R motifs in a broad collection of substrates, including human histone peptides and RG-rich peptides. In addition, the activity of the PRMT-7 enzyme is dependent on temperature, the presence of metal ions, and the reducing agent dithiothreitol. *C. elegans* PRMT-7 has a substrate specificity and a substrate preference different from those of mammalian PRMT7, and the available X-ray crystal structures of the PRMT7 orthologs show differences in active site architecture. *C. elegans* PRMT-9, on the other hand, produces symmetric dimethylarginine and MMA on SFTB-2, the conserved *C. elegans* ortholog of human RNA splicing factor SF3B2, indicating a possible role in the regulation of nematode splicing. In contrast to PRMT-7, *C. elegans* PRMT-9 appears to be biochemically indistinguishable from its human ortholog.



Protein arginine methylation is a posttranslational modification that is widely distributed in vertebrates, invertebrates, plants, and fungi.^{1,2} Members of the mammalian protein arginine methyltransferase (PRMT) family modify arginine residues by introducing a methyl group onto the terminal guanidino groups, generating either exclusively monomethylarginine (MMA) (type III, PRMT7), MMA and symmetric dimethylarginine (SDMA) (type II, PRMT5 and PRMT9), or MMA and asymmetric dimethylarginine (ADMA) (type I, PRMT1–4, -6, and -8).^{3,4} This family of enzymes has roles in the epigenetic regulation of transcription,⁵ mRNA splicing,⁵ DNA damage response,⁶ carcinogenesis,^{3,7,8} and signaling.⁹ Many of these enzymes have multiple functions and roles, and changes in their expression have been implicated in tumorigenesis and disease.^{3,4,7,8}

The majority of PRMTs have been well characterized, and their physiological substrates have been identified. PRMT7 and PRMT9, however, have received much less attention. While PRMT9 appears to have a major role in the regulation of alternative splicing,⁵ for PRMT7 the situation is more complex. Recent studies have shown that PRMT7 is involved in preserving satellite cell regenerative capacity,¹⁰ regulation of germinal center formation,¹¹ induction of the epithelial-to-mesenchymal transition,¹² and bone development,¹³ in addition to roles in cancer such as the promotion of breast cancer metastasis through MMP9 expression.¹⁴ However, it has not

yet been possible to connect these phenomena to the specific enzymatic action of PRMT7. In addition, there is evidence that crosstalk of PRMT7 with other PRMTs can modify the epigenetic code. For instance, the knockdown of PRMT7 in mammalian cells has been shown to decrease the level of SDMA formation at arginine 3 in histone H2A and H4 in chromatin associated with specific genes^{10,15,16} in spite of the fact that PRMT7 does not catalyze SDMA formation.^{3,17,18} In addition, there is evidence of the association of increased levels of arginine 3 SDMA in histone H4 at specific genes with overexpression of PRMT7.¹¹ It thus appears that PRMT7 may function in conjunction with other PRMTs.

Because of the complexity of mammalian systems, PRMT7 from lower eukaryotes, including *Trypanosoma brucei*^{19–22} and *Caenorhabditis elegans*,^{23,24} has been studied. However, the trypanosome enzyme has a preference for RG-rich proteins such as RBP16,^{19,21,22} while mammalian PRMT7 has a preference for R-X-R motifs in basic sequence contexts such as those found in histone H2B.^{17,18} A crystal structure is available for *C. elegans* PRMT-7 that is similar to that determined for *Mus musculus* PRMT7, with both enzymes containing a single chain with two ancestrally duplicated

Received: March 28, 2017

Revised: April 20, 2017

Published: April 25, 2017

methyltransferase domains stabilized by a zinc ion.^{24,25} We note that while the *T. brucei* ortholog is named PRMT7, its structure is divergent from those of the *C. elegans* and mouse enzymes, lacking the duplicated domain characteristic of plant, invertebrate, and vertebrate PRMT7 members.^{1,2,19} A previous attempt to characterize *C. elegans* PRMT-7 (then named PRMT-2) *in vitro* was unsuccessful because it was enzymatically inactive under the conditions tested.²³ The authors of this same study²³ designated *C. elegans* PRMT-9 (then named PRMT-3) as a human PRMT7 ortholog, while more recent studies have shown it is more closely related to human PRMT9.^{1,26}

Given the similarity of the *C. elegans* and mouse PRMT7 enzymes, and the difficulty of determining the mechanism of PRMT7 action in vertebrates, we sought to characterize the *C. elegans* ortholog. We have now been able to show that although the *C. elegans* enzyme is similar to the previously studied mammalian enzyme in its ability to form only MMA marks,^{17,18,27} the nematode enzyme has distinct specificity for recognizing arginine residues in peptides and proteins, suggesting distinct functions.

Mammalian PRMT9 had also been challenging to characterize. None of the previously characterized PRMT methyl acceptors were recognized by human PRMT9.²⁶ However, it was possible to identify a complex with splicing factors SF3B2 and SF3B4 and to show that PRMT9 methylates SF3B2 to produce MMA and SDMA.^{5,26} SF3B2 may be the only major substrate of this enzyme, and its modification appears to play a role in the regulation of alternative splicing.^{5,26}

While *C. elegans* has no orthologs of mammalian PRMT2–4, -6, or -8, there is an ortholog of PRMT9.^{1,2} In the previous study, this ortholog was shown to methylate recombinant histone H2A, producing only MMA.²³ Thus, we were interested in characterizing this *C. elegans* PRMT-9 to establish whether it plays a role similar to that of mammalian PRMT9. We now show that *C. elegans* PRMT-9 is a type II enzyme that produces MMA and SDMA, but it also appears to be highly specific for the *C. elegans* ortholog of the SF3B2 splicing factor, SFTB-2.

MATERIALS AND METHODS

Bacterial Protein Expression and Purification. *C. elegans* PRMT-7 cDNA and PRMT-9 cDNA were cloned into a pGEX-6P-1 plasmid²³ and were a kind gift from A. Fukamizu. The proteins were expressed in BL21 DE3 cells (Invitrogen); *C. elegans* GST-PRMT-7 and -9 enzymes were purified using the conditions described in ref 26, but with induction for 20 h at 18 °C using 0.13 mM isopropyl D-thiogalactopyranoside (IPTG) for GST-PRMT-7 and 1 mM IPTG for 20 h at 18 °C for both GST-PRMT-9 wild type and A391H mutant. The enzymes were purified as described and dialyzed overnight into 100 mM Tris-HCl and 100 mM NaCl (pH 7.5). *Homo sapiens* GST-PRMT7,¹⁸ GST-PRMT9,²⁶ and GST-SF3B2^{5,26} were expressed and purified as described previously. GST-GAR was expressed using 0.4 mM IPTG for 20 h at 18 °C and purified using a process similar to that used for the other proteins but was dialyzed into 50 mM HEPES, 120 mM NaCl, and 1 mM DTT (pH 8.0). The amino acid sequence for SFTB-2 (*C. elegans* SF3B2) was synthesized by GenScript, Inc., and cloned into a pGEX-6p-1 vector. GST-tagged SFTB-2 was purified using a process similar to that used for human SF3B2 and dialyzed overnight into 10 mM Na₂HPO₄, 2 mM KH₂PO₄, 137 mM NaCl, 2.7 mM KCl, and 1 mM DTT (pH 7.4).

Each of these plasmids encodes the full sequence of the *Schistosoma japonicum* glutathione S-transferase (UniProt entry P08515) followed by linker regions of SDLEVLFGQPLGSGIP (*C. elegans* PRMT-7), SDLEVLFGQPLGSPEFP (*C. elegans* PRMT-9), SDLQVLFQGPL (*C. elegans* SFTB-2), SDLVPRG-SST (human PRMT7), and SDLVPRGS (GST-GAR) followed by the full amino acid sequences of either *C. elegans* PRMT-7 (UniProt entry Q9XW42), *C. elegans* PRMT-9 (UniProt entry O02325), human PRMT7 (UniProt entry Q9NVM4), human PRMT9 (UniProt entry Q6P2P2), *C. elegans* SFTB-2 (UniProt entry O16997), human SF3B2 (UniProt entry Q13435), or amino acids 1–145 of human fibrillarin (UniProt entry P22087), the latter of which has K2E and A145V substitutions. The GST-GAR fusion protein was previously erroneously reported to include human fibrillarin residues 1–148.^{18,28} Purified proteins were analyzed via sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE) and then quantified using 10% trichloroacetic acid precipitation for a Lowry assay.


Peptide Synthesis. All H2B peptides used in these studies listed in Table S1 were synthesized by GenScript, Inc., and further validated by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. Other peptides were gifts from their respective sources (see Table S1).

In Vitro Methylation Reactions. Methylation reactions included 2 µg of enzyme and potential methyl-accepting protein substrate (5 µg) or peptide substrates at a final concentration of 12.5 µM. Mixtures were incubated at the indicated temperatures using a buffer of 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) unless otherwise indicated and 0.7 µM S-adenosyl-L-[methyl-³H]methionine {³H-AdoMet, PerkinElmer Life Sciences, 82.7 Ci/mmol, 0.55 mCi/mL in 10 mM H₂SO₄/EtOH [9:1 (v/v)]} in a final reaction volume of 60 µL. Given a counting efficiency of 50%, 1 fmol of radiolabeled methyl groups corresponds to 91 cpm.

Amino Acid Analysis of Substrates Using High-Resolution Cation Exchange Chromatography. Reactions with peptide substrates were stopped by the addition of 3 µL of 25% trichloroacetic acid, and peptides were purified using OMIX C18 Zip-Tip pipet tips (Agilent Technologies). The reaction mixtures were then subjected to vacuum centrifugation to remove the Zip-Tip elution buffer consisting of H₂O and acetonitrile (50:50). Protein substrate reactions were stopped by the addition of an equal volume of 25% (w/v) trichloroacetic acid followed by centrifugation to give the protein pellet. Methylated peptides and proteins were subsequently acid hydrolyzed, mixed with nonradioactive methylated arginine derivatives, and separated and analyzed using high-resolution cation exchange chromatography as described in ref 5.

Detection of Methylated Substrates after SDS–PAGE. Protein substrate methylation reactions were stopped by the addition of 0.2 volume of 5× SDS sample loading buffer and run on a 12.6% polyacrylamide Tris-glycine gel. The gel was stained using Coomassie Blue, and after being destained overnight, gels were incubated in EN³HANCE (PerkinElmer Life Sciences, catalog no. 6NE9701) autoradiography reagent, washed with water for 30 min, and then vacuum-dried. Dried gels were then placed in a cassette and exposed to autoradiography film (Denville Scientific, catalog no. E3012) at –80 °C for the amount of time indicated in the figure legends. Densitometry of the bands was performed using ImageJ.

A



B




Figure 1. Evolutionary conservation of PRMT7 across the various kingdoms of life. (A) Phylogenetic tree based on human PRMT7. UniProt entries of representative orthologs from Animalia, Plantae, Archaeplastida, Fungi, Bacteria, and Excavata are shown with their *E* values based on a protein BLAST search against the human species.²⁹ The phylogenetic tree was constructed after amino acid sequences were aligned using MUSCLE in MEGA6 software as described in ref 26. Each ortholog sequence was then subjected to protein BLAST against the human protein database. All proteins were mutual best hits with the exception of those species marked with an asterisk. The number shown next to each branch is the percent of replicate trees with the same clustering in 500 bootstrap test replicates. The scale bar indicates the fraction of amino acid differences for each entry. *H. sapiens* PRMT7 is boxed in dark red, and *C. elegans* PRMT-7 is boxed in red. (B) Partial sequence alignment of the major motifs in PRMT7 orthologs across vertebrates, invertebrates, plants, and excavata (*T. brucei*) using the EMBL-EBI Clustal Omega Multiple Sequence Alignment software.²⁹ The major distinguishing motifs of protein arginine methyltransferases are boxed in black, including motif I, post motif I, motif II, the double E loop, and the THW loop. Red letters indicate identity and blue letters similar amino acid properties. The secondary structure is indicated on the bottom with β -strands colored yellow and α -helices colored magenta based on the structure of *M. musculus* PRMT7 (Protein Data Bank entry 4C4A²⁵). The *C. elegans* PRMT-7 sequence is boxed in dark gray.




Figure 2. *C. elegans* PRMT-7 produces MMA, and its activity is dependent on temperature and metal ion. (A) Amino acid analysis of ^3H methylation reactions of *C. elegans* GST-PRMT-7 ($2\ \mu\text{g}$) with GST-GAR ($5\ \mu\text{g}$) after a 20 h reaction at $30\ ^\circ\text{C}$ (left) or $25\ ^\circ\text{C}$ (middle and right) in either Tris-EDTA buffer [100 mM Tris-HCl, 100 mM NaCl, and 2 mM EDTA (pH 8.0)] (top) or 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) (bottom) as indicated in [Materials and Methods](#). The final concentration of DTT in these reactions (from the GST-GAR preparation) was 0.16 mM. After trichloroacetic acid precipitation and acid hydrolysis, $1\ \mu\text{mol}$ of each nonradiolabeled methylated arginine standard (ADMA, SDMA, and MMA) was added to the hydrolyzed pellet and amino acid analysis was performed as described in [Materials and Methods](#). Dashed black lines indicate ninhydrin absorbance of the methylarginine standards with the peaks of ADMA, SDMA, and MMA identified using $50\ \mu\text{L}$ of each fraction. Red lines indicate the elution of the radiolabeled methylated amino acids from the hydrolysates of the reactions. Radioactivity from $950\ \mu\text{L}$ aliquots of each fraction is given as the average of three 5 min counting cycles using liquid scintillation counting. Radioactive methylated amino acids elute approximately 1 min before the nonradiolabeled standard because of the tritium isotope effect.³⁶ The right panels show an expanded view of the data from the middle columns at $25\ ^\circ\text{C}$ to demonstrate that the enzyme produces MMA only under both buffer conditions. These reactions were replicated three independent times. The asterisk indicates the peak eluting prior to the position expected for ^3H -ADMA; this peak is not consistently observed in the replicates. (B) PRMT-7 enzyme activity is dependent on the presence of DTT. The enzyme was reacted with $12.5\ \mu\text{M}$ pSmD3 (top) or pRpI3 (bottom) peptides at $25\ ^\circ\text{C}$ in 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) (left) or at $15\ ^\circ\text{C}$ in the same buffer with the addition of 1 mM DTT. Results are presented as shown in panel A; the blue line in the right-hand panels indicates radioactivity for incubations to which no peptide substrate was added. These reactions were replicated twice with no DTT and once with DTT.

RESULTS

PRMT7 Is Widely Conserved in Vertebrates, Invertebrates, and Plants. The human PRMT7 sequence was analyzed by protein BLAST,²⁹ and the representative orthologs were compiled into a phylogenetic tree (Figure 1A). We found that the major PRMT structural motifs, including AdoMet-binding post motif I, the substrate-binding double E loop, and the THW loop, are well-conserved (Figure 1B). PRMT7 is widely distributed across vertebrates and invertebrates, including mollusks, insects, tunicates, and nematodes, as well

as in higher plants (Figure 1A and ref 1). Significantly, the *C. elegans* ortholog is 32% identical to the human enzyme over the full polypeptide length and has an expected value of 3×10^{-96} . Fungi appear to lack a PRMT7 ortholog, with the closest relatives being more similar to PRMT1. Bacteria also seem to largely lack orthologs to PRMT7. The closest bacterial sequence homologues in *Streptococcus pneumoniae* and *Pseudanabaena* sp. are not mutual best hits to PRMT7, although the *Rhizobium leguminosarum* and *Geobacter sulfurreducens* species are most closely related to PRMT7. In addition,




Figure 3. PRMT-7 has a temperature dependence consistent with *C. elegans* physiology. Methylation reaction mixtures consisting of 5 μg of GST-GAR substrate and 2 μg of enzyme (*C. elegans* GST-PRMT-7 or *H. sapiens* GST-PRMT7) were incubated for 20 h at the indicated temperatures in reaction buffer containing 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) with 0.7 μM [methyl- ^3H]AdoMet in a final reaction volume of 60 μL . The final concentration of DTT in these reactions (from the GST-GAR preparation) was 0.16 mM. Reactions were quenched by the addition of SDS sample loading buffer and polypeptides separated by 12.6% Tris-glycine SDS-PAGE followed by autoradiography as described in Materials and Methods. The dried gel was then exposed to Denville E3012 autoradiography film for 21 days at -80°C . Molecular weight positions are shown from approximately 2 μg of unstained SDS-PAGE broad range marker (Bio-Rad, catalog no. 161-0317). This full experiment was independently replicated, and a further third replicate using only the *C. elegans* enzyme was also performed. Densitometry was performed using ImageJ on scanned images of the films and quantified as relative density and normalized to the highest peak of GST-GAR radioactivity in each individual film (*C. elegans*, red symbols; human, blue symbols; different symbols used to indicate each replicate experiment). Lines are drawn for the averaged normalized values. The optimal growth temperatures for *C. elegans*³¹ and humans are indicated. Lower bands shown on the fluorography gel are nonspecific and not dependent on enzyme activity.

the type III (MMA forming) trypanosome *T. brucei* PRMT7^{19,21} is most similar in sequence to human PRMT9 and is not a mutual best hit for human PRMT7. Importantly, *T. brucei* PRMT7 has a substrate specificity distinct from those of




Figure 4. *C. elegans* PRMT-7 has an R-X-R substrate specificity similar to that of the human PRMT7 enzyme. (A) Amino acid analysis of reaction mixtures consisting of 2 μg of human GST-PRMT7 or *C. elegans* PRMT-7 enzymes, reacted with 12.5 μM synthetic human histone H2B (23–37) peptide^{17,18} or a synthetic peptide containing the equivalent sequence from *C. elegans* histone H2B (20–34) at 15 $^\circ\text{C}$ as described in Materials and Methods, in 50 mM potassium HEPES buffer and 10 mM NaCl (pH 8.0) containing 1 mM DTT for 20 h. Reactions were repeated three times for human histone H2B (23–37) and twice for *C. elegans* H2B (20–34) at this temperature and also replicated at 25 $^\circ\text{C}$ once. (B) Amino acid analysis of reaction mixtures containing 2 μg of *C. elegans* GST-PRMT-7 enzyme reacted with 12.5 μM synthetic human histone H2B peptides (wild type, top; R29K, R31K, and R33K bottom) for 20 h, using the same conditions described above for panel A. These reactions were single replicates. (C) Amino acid analysis of reaction mixtures containing 2 μg of *C. elegans* or human PRMT-7/7 enzymes, reacted with 12.5 μM synthetic peptide RGR-1 (a sequence containing an R-X-R sequence where X = G) at 25 $^\circ\text{C}$ (left) or 15 $^\circ\text{C}$ (right), under the same reaction conditions as described for panel A. Reactions were replicated once, and the reaction of *C. elegans* with RGR-1 was replicated two additional times.

both human PRMT7 and PRMT9, preferring RG-rich proteins.^{5,17–19,21,22,26}

The major conserved PRMT motifs are shown for selected species in Figure 1B. While all of the enzymes have similar motifs common to all seven β -strand methyltransferases (motif I, post motif I, and motif II), the *T. brucei* PRMT7 deviates significantly from the other proteins in the PRMT-specific




Figure 5. *C. elegans* PRMT-7 has a preference for R-X-R motifs in various peptide substrates. (A) *C. elegans* PRMT-7 (2 μg) was reacted with 12.5 μM various synthetic human histone H4 peptides (1–21 wild type, 1–21 R3K, 1–16 wild type, and 14–22) at 15 °C in 50 mM potassium HEPES and 10 mM NaCl (pH 8.0) with 1 mM DTT for 20 h. Amino acid analysis was performed as described above and in [Materials and Methods](#). Reactions were replicated twice. (B) *C. elegans* PRMT-7 (2 μg) was reacted with various synthetic peptides [pSmD3, pRpl3, pR1, histone H3 (1–7), and RGR-1] at 12.5 μM under the conditions described for panel A. Reactions were single replicates. The reactions in the top panels for pSmD3 and pRpl3 are the same panels used in the right column of [Figure 2B](#).

double E loop and the THW loop. When compared to the human PRMT7, the *C. elegans* protein has an identical DHW sequence in the THW loop as well as an identical double E loop with the exception of the replacement of a single leucine residue with a valine. These results indicate that PRMT7 is evolutionarily conserved with a marked presence in vertebrates, invertebrates, and plants, but not in lower organisms.

***C. elegans* PRMT-7 Produces Monomethylarginine.** When we assayed the *C. elegans* PRMT-7 methyltransferase activity under conditions used in the previous biochemical study²³ (30 °C with an EDTA-containing buffer), we also found no activity ([Figure 2A](#), top left panel). However, when the incubation temperature is decreased to one more favorable for *C. elegans* growth (25 °C), some enzymatic activity producing MMA was seen ([Figure 2A](#), top middle and right

panels). When we switched to a buffer containing no metal ion chelator, some activity was seen at 30 °C ([Figure 2A](#), bottom left panel) while much higher activity was seen at 25 °C ([Figure 2A](#), bottom middle and right panel). Previous work had demonstrated the partial loss of activity of the mammalian PRMT7 enzyme with EDTA.¹⁸ A zinc ion is present in a zinc finger motif in the mouse and *C. elegans* PRMT7/PRMT-7 structures located some distance from the active site, suggesting that it may play a role in stabilizing an active conformation of the enzyme.^{24,25} Thus, it appears that EDTA may inhibit the activity by extracting this zinc ion. In addition, the *C. elegans* PRMT-7 enzyme activity appears to be dependent on the presence of the reducing agent dithiothreitol (DTT), as was previously observed with the mammalian enzyme.²⁷ *In vitro* methylation reaction mixtures with two synthetic peptides




Figure 6. *C. elegans* PRMT-7 has a substrate specificity distinct from that of the mammalian ortholog for mammalian histones. Reaction mixtures consisting of 5 μ g of substrate [recombinant human histones from New England BioLabs (H2A, M2502S; H2B, M2505S; H3.3, M2507S; H4, M2504S)] were reacted with 2 μ g of *C. elegans* GST-PRMT-7 or human GST-PRMT7 at 15 or 25 $^{\circ}$ C as described in [Materials and Methods](#). Reactions were stopped by the addition of sample loading buffer and polypeptides separated as described in [Materials and Methods](#). The dried gel was then exposed to autoradiography film for 3 days at -80° C. Molecular weight positions are shown from approximately 2 μ g of unstained SDS-PAGE broad range marker as in [Figure 3](#). The results shown here were replicated three times for *C. elegans* PRMT-7 and two times for human PRMT7. Radiolabeled bands migrating more rapidly than histone H2B in the lane with human PRMT7 appear to be proteolytic fragments.

containing R-X-R and similar sequences (pSmD3 and pRpl3) showed no activity at 25 $^{\circ}$ C without DTT ([Figure 2B](#), left panels) but showed significant MMA-producing activity when DTT was added to the reaction buffer ([Figure 2B](#), right panels) at 15 $^{\circ}$ C.

On the basis of reports that the expression level of PRMT7 can affect SDMA levels in mammalian histones,^{11,12,15,16,30} we also asked if the active *C. elegans* PRMT-7 enzyme could catalyze dimethylation of arginine species. However, as shown in the expanded view of the amino acid analysis shown in [Figure 2A](#) (right panels), we find no dimethylated product consistent with SDMA or ADMA is formed under conditions where 0.4% of the MMA product would have been detected. Thus, *C. elegans* PRMT-7 produces only MMA and is thus a type III enzyme.

***C. elegans* PRMT-7 Is Most Active at or below Optimal Growth Temperatures.** In [Figure 2](#), we noted the enzyme was much more active at 25 $^{\circ}$ C than at 30 $^{\circ}$ C. *C. elegans* is able to grow at a variety of temperatures ranging from 12 $^{\circ}$ C (slowest-growing) to 25 $^{\circ}$ C (fastest-growing), with most studies performed at 20 $^{\circ}$ C.³¹ Continual incubation at $>25^{\circ}$ C leads to sterility.^{31,32} We thus performed *in vitro* methylation reactions with *C. elegans* PRMT-7 and GST-GAR over a wider range of temperatures, from 0 to 42 $^{\circ}$ C ([Figure 3](#)). After SDS-PAGE and fluorography, the methylation signal was quantified using densitometry and normalized. Fluorography revealed an activity maximum at 15 $^{\circ}$ C, with the enzyme still slightly active at the higher end of the physiological temperature range for *C. elegans* (20 and 25 $^{\circ}$ C). *In vitro* methylation reactions of human PRMT7 were performed as a comparison, and a similar but narrower dependence of activity on temperature is observed ([Figure 3](#)). The temperature dependence of the human enzyme observed here with a maximal activity at 15 $^{\circ}$ C and little or no activity at 37 $^{\circ}$ C is similar to what was reported previously.¹⁸

***C. elegans* PRMT-7 Recognizes a Wider Variety of Substrates Than Human PRMT7 Does.** Once reaction conditions were optimized for activity, we tested a variety of

protein and peptide substrates to compare the methyl-accepting activity with peptide and protein substrates, including those recognized by human PRMT7. Because both human and mouse PRMT7 were shown to have a strong preference for R-X-R motifs flanked by basic residues in histone H2B peptides,^{17,18} we decided to test these peptides with *C. elegans* PRMT-7. Amino acid analysis showed that *C. elegans* PRMT-7 can methylate synthetic human histone H2B peptides containing residues 23–37 ([Figure 4A](#), bottom left panel). To determine whether the *C. elegans* enzyme better recognizes *C. elegans* amino acid sequences than human sequences, we performed the same experiment with the *C. elegans* histone H2B (20–34) peptide. Surprisingly, we found that the *C. elegans* enzyme recognized the human peptide better than the *C. elegans* peptide ([Figure 4A](#), bottom). We compared this activity to that of human PRMT7 with the human peptide and the *C. elegans* peptide ([Figure 4A](#), top panel) and found that the human enzyme did not methylate the *C. elegans* H2B peptide sequence. These results suggest that, at least *in vitro*, histone H2B may not be a major substrate of PRMT-7 in *C. elegans*.

Because *C. elegans* PRMT-7 was able to methylate the human histone H2B peptide, we then tested our collection of single R-to-K mutants of this peptide ([Figure 4B](#)). When it was reacted with the wild-type H2B peptide, we see similar results as in [Figure 4A](#), but the removal of a single arginine from the peptide greatly decreased the activity ([Figure 4B](#), bottom panels). The histone H2B R29K peptide reduced methylation counts to a level that was $\sim 40\%$ of that of the wild-type peptide sequence. The R31K and R33K mutant peptides further decreased the activity to ~ 10 and $\sim 22\%$, respectively, signifying the importance of arginine residues in a sequential R-X-R motif. Because our data showed that PRMT-7 prefers an R-X-R motif in histone H2B, we tested another synthetic peptide containing an R-X-R sequence, where X is a glycine, a sequence found in many GAR motif proteins. We tested the *C. elegans* and human enzymes with this RGR-1 peptide at 25 and 15 $^{\circ}$ C and found that the *C. elegans* enzyme weakly methylated this peptide at both temperatures ([Figure 4C](#), top row), while the human




Figure 7. Active site architecture of *C. elegans*, trypanosome, and mammalian PRMT7s. (A) Pymol surface and cartoon representation of the active site of *C. elegans* PRMT-7 (Protein Data Bank entry 3XOD²⁴). Flanking glutamate double E loop residues are colored cyan (E140 and E149); internal loop acidic residues are colored purple (D143 and E145), and a THW loop residue is colored magenta (H300). The residue colored orange (F33) protrudes from a helix to form a cavity with E149. The distance from the side chain carboxyl carbon of E140 to the side chain carboxyl carbon of E149 is 7.6 Å, and the distance from the side chain carboxyl carbon of E149 to the closest side chain carbon atom in F33 is 5.2 Å. S-Adenosylhomocysteine (SAH) is shown in CPK coloring. (B) Pymol surface and cartoon representation of the active site of *T. brucei* PRMT7 (Protein Data Bank entry 4M38²⁰), highlighting residues corresponding to those shown in panel A. Flanking double E loop residues E172 and E181 are colored cyan; G175 and M177 (residues corresponding to D143 and E145, respectively, in *C. elegans*) are colored purple, and the Q329 THW loop residue is colored magenta. The residue colored orange (F71) corresponds to F33 in the *C. elegans* structure and also protrudes from a helix to form a cavity with E181. The distances between the flanking double E residues and E181 and F71 are given as in panel A. The arginine 3 residue of the cocrystallized histone H4 peptide is colored red. (C) Pymol surface and cartoon representation of the active site of *M. musculus* PRMT7 (Protein Data Bank entry 4C4A²⁵), highlighting residues corresponding to those shown in panels A and B. The distances between the flanking double E residues and E153 and S34 are given as in panels A and B. We note that the S34 side chain oxygen is shown in two conformations.

PRMT7 enzyme showed much less activity (Figure 4C, bottom row), indicating a slightly broader specificity for the *C. elegans* enzyme.

Once we saw the differences in substrate specificity of the *C. elegans* PRMT-7 enzyme when compared to that of the previously characterized human enzyme, we tested it against another panel of synthetic peptides with various sequences

containing pairs of arginine residues in R-X-R motifs as well as single-arginine sequences. Upon incubation with synthetic histone H4 peptides (1–21 wild type, 1–21 R3K, 1–16, and 14–22), we saw greatly reduced activity when the R-X-R sequence at the C-terminus of the peptide was removed (Figure 5A). This peptide contained the sequence of the *C. elegans* histone, one identical to that of the human peptide. The histone H4 (1–21) R3K peptide gave just a slight decrease in methylation activity [20% decrease (Figure 5A, top right panel)], but removal of the last five residues in the peptide of residues 1–16 caused an almost 70% decrease in activity (Figure 5A, bottom left panel). This signified that the PRMT-7 enzyme primarily methylates the R-X-R sequence but could also recognize and methylate a single arginine. Reacting the enzyme with a small peptide containing only the R-X-R sequence of H4 (14–22) yielded almost no activity, indicating there might be a secondary site of contact and binding for the peptide to be correctly situated in the active site (Figure 5A, bottom right panel).

Other peptides containing R-G-R and peptides containing a single arginine residue were reacted with the *C. elegans* enzyme (Figure 5B). In a longer peptide sequence containing two pairs of R-G-R sequences [pSmD3 (Figure 5B, top left panel)], we saw very high methylation activity. In a reaction mixture with pRpl3 peptide (Figure 5B, top middle panel) that contains three more widely spaced arginine residues, we see an activity that is almost double that of that seen with the RGR-1 peptide (Figure 5B, bottom right panel). *C. elegans* PRMT-7 was not able to methylate smaller peptides, such as pR1 (containing only one R flanked by G) or a peptide containing the first seven residues from histone H3 (Figure 5B, bottom panels).

We then tested full-length histones as substrates of both the human and *C. elegans* PRMT7 enzymes. We methylated recombinant human histone proteins at either 15 or 25 °C and separated the polypeptides via SDS–PAGE. At both temperatures, we found that the *C. elegans* PRMT-7 enzyme was active in methylating histones H2A, H2B, and H3, but only weakly methylated H4 (Figure 6). This was a surprising result because the enzyme could methylate H4 peptides (Figure 5A). We note that the sequence of histone H4 in humans and *C. elegans* is identical with the exception of one residue in the central domain. Human PRMT7, on the other hand, was very specific for histone H2B as noted previously,¹⁷ indicating a distinct difference in substrate specificity between the two enzymes.

Crystal Structures of PRMT7 Orthologs Unveil a Distinct Pocket in the Active Site Potentially Contributing to Substrate Specificity. Although the characterized mammalian, trypanosome, and *C. elegans* PRMT7 homologues all produce MMA, there are significant differences in their substrate specificity (this study and refs 18 and 22). We therefore examined the active site architecture of the available PRMT7 crystal structures of each of these enzymes. The substrate arginine residue is generally bound between the two glutamate residues in the double E loop.^{3,4,22} The distance between the side chain carboxyls of the two glutamates in the double E loop is relatively similar in the *C. elegans* and *T. brucei* enzymes (7.6 and 7.2 Å, respectively), compared to a larger distance (8.7 Å) in the *M. musculus* enzyme (Figure 7). In addition, in PRMT7 from *C. elegans* and *T. brucei*, there is a phenylalanine residue that protrudes into the active site (F33 and F71) and may limit the size of a methyl-accepting sequence. The distance between the side chain carboxyl carbon





Figure 8. *C. elegans* PRMT-9 symmetrically dimethylates SFTB-2, the *C. elegans* ortholog of the mammalian SF3B2. (A) Amino acid analysis was performed as described in the legend of Figure 2 on methylation reaction mixtures consisting of 5 μg of substrate (*C. elegans* GST-tagged SFTB-2 fragment 99–248 or *H. sapiens* GST-tagged SF3B2 fragment 401–550) and approximately 2 μg of enzyme (*C. elegans* GST-tagged PRMT-9 enzyme or *H. sapiens* GST-tagged PRMT9 enzyme) that were incubated for 20 h at 25 °C (nematode enzyme) or 37 °C (human enzyme) in a reaction buffer consisting of 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) with a final concentration of 0.7 μM [methyl-³H]AdoMet in a volume of 60 μL. The reactions were quenched by the addition of a final trichloroacetic acid concentration of 12.5% and mixtures acid-hydrolyzed as described in Materials and Methods. The asterisk indicates a radioactive peak eluting prior to the expected position of ³H-ADMA. The experiment was replicated twice. (B) Sequence alignment of *C. elegans* SFTB-2 (UniProt entry O16997) residues 99–248 and human SF3B2 (UniProt entry Q13435, bottom). Red letters indicate identity, and blue letters indicate similar amino acid properties. The bolded R and black arrow indicate the methylation site for the enzyme.

of the second glutamate residue of the double E loop and the closest side chain carbon of the phenylalanine residue is 5.2 and 5.8 Å in *C. elegans* and *T. brucei*, respectively (Figure 7A,B). However, the corresponding residue in *M. musculus* PRMT7 is a serine residue whose β-carbon is much farther (8.5 Å) from the glutamate residue (Figure 7C), potentially creating a pocket that could accommodate arginine residues flanked by residues with bulkier side chains such as lysine residues. We also note that the two acidic residues in the interior of the double E loop (colored purple in Figure 7) are more exposed to the surface in

M. musculus (Figure 7C) than the corresponding residues in the *C. elegans* or *T. brucei* enzymes. These results provide insight into the broader substrate specificity of the non-mammalian PRMTs.

The *C. elegans* PRMT-9 Enzyme Methylates RNA Splicing Factor SFTB-2/SF3B2 and Forms SDMA. We previously showed a phylogenetic tree demonstrating a *C. elegans* ortholog of PRMT9 present in higher organisms;²⁶ this ortholog was originally named PRMT-3 but is now designated PRMT-9.^{1,23,26} In addition, *C. elegans* also contains an ortholog




Figure 9. *C. elegans* PRMT-9 methylates SF3B2 at the same R508 site that is methylated by the mammalian PRMT9 enzyme. (A) Amino acid analysis of a methylation reaction mixture consisting of 2 μ g of *C. elegans* GST-PRMT-9 enzyme with 5 μ g of human GST-SF3B2 (401–550) wild type at 25 °C for 20 h in either 100 mM Tris-EDTA buffer (pH 8.0) (top) or 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) (bottom). Reactions were quenched by the addition of a final trichloroacetic acid concentration of 12.5% and mixtures acid-hydrolyzed as described above. Reactions were duplicated twice. (B) Methylation reactions of *C. elegans* PRMT-9 reacted with the human GST-SF3B2 (401–550) R508K mutant fragment, under the same reaction conditions described for panel A at 25 °C for 20 h. Reactions were duplicated twice.

of mammalian splicing factor SF3B2, now designated SFTB-2. SF3B2 is the only identified substrate of mammalian PRMT9.^{5,26} To determine if *C. elegans* PRMT-9 plays a similar physiological role by methylating SFTB-2, we analyzed the products of both human and *C. elegans* PRMT-9 enzymes with the methylatable fragment F3 of both splicing factors, corresponding to human residues 401–550 and *C. elegans* residues 99–248 (Figure 8A). When the *C. elegans* enzyme was incubated with the *C. elegans* SFTB-2 and mammalian SF3B2 fragments, we detected both MMA and SDMA formation (Figure 8A, top panels). In a control experiment, we also detected MMA and SDMA formation with the human enzyme and substrate (Figure 8A, bottom right panel). However, when we reacted the human enzyme with the *C. elegans* SFTB-2 polypeptide, we found only a small MMA peak; no SDMA was detected (Figure 8A, bottom left panel). SF3B2 and SFTB-2 are similar in sequence in the conserved region surrounding the R508 methylation site in the human protein^{5,26} but diverge in the N-terminal region (Figure 8B).

To ask if the *C. elegans* enzyme was specific for the same site, we reacted *C. elegans* PRMT-9 with the wild type and R508K mutant of the human SF3B2 fragment (Figure 9). We observed that methylation was abolished in the fragment with the R508K mutation (Figure 10B, top and bottom right panels), as

compared to the case for the wild-type SF3B2 fragment (Figure 10A, top and bottom left panels), suggesting that the mammalian and *C. elegans* enzymes share the same strict specificity for this arginine residue in the splicing factor.

***C. elegans* PRMT-9 Does Not Recognize Non-SFTB-2/SF3B2 Substrates.** Because most PRMTs recognize a variety of substrates, we tested multiple known substrates with the *C. elegans* enzyme, including GST-GAR, RNA binding protein Rbp16, and recombinant histones H2A and H2B (Figure 10). Amino acid analysis of reactions using the same conditions as described by Takahashi et al.,²³ the first study to detect MMA activity with this PRMT-9 enzyme, showed that it did not appreciably methylate these substrates (Figure 10A, top right panel). We did, however, observe a very low MMA activity upon the methylation of recombinant histone H2A, consistent with the previous result.²³ We also analyzed these reaction mixtures, along with corresponding mixtures with the human PRMT9 enzyme, using SDS-PAGE fluorography, and observed that both PRMT9 enzymes methylated only their corresponding SF3B2 substrate fragments (Figure 10B).

Temperature Dependence of *C. elegans* PRMT-9. We next compared the activity at various temperatures of both the human and nematode PRMT9 enzymes with the human SF3B2 substrate using SDS-PAGE fluorographic analysis (Figure 11).




Figure 10. Substrate specificity of *C. elegans* PRMT-9. (A) Methylation reactions for amino acid analysis were set up using 2 μ g of *C. elegans* PRMT-9 with 5 μ g of various substrates [GST-GAR, top left; GST-SF3B2 (401–550), top right; recombinant human histone H2A, bottom left; recombinant human histone H2B, bottom right], in 100 mM Tris-EDTA buffer (pH 8.0) for 6 h at 30 °C, under conditions defined by Takahashi et al.²³ These reactions using these conditions were single replicates. (B) *In vitro* methylation reaction mixtures consisting of 2 μ g of *C. elegans* PRMT-9 or human PRMT9 enzyme were reacted with 5 μ g of substrate [GST-GAR, His-Rbp16, or GST-SFTB-2 (*C. elegans* fragment 99–248) or GST-SF3B2 (human fragment 401–550)] for 20 h at 25 °C (*C. elegans*) or 37 °C (human) in 50 mM potassium HEPES, 10 mM NaCl, and 1 mM DTT (pH 8.2) with 0.7 μ M [*methyl*-³H]AdoMet. Sample loading buffer was added to stop the reactions, and polypeptides were separated using SDS-PAGE, using the same methods described in the legend of Figure 3. The gel was then dried and exposed to film for 7 days at –80 °C. The Coomassie-stained gel is shown in the top panel, and the fluorographic exposure is shown in the bottom panel. The molecular weight standards are shown as described in the legend of Figure 3. This experiment was replicated independently using amino acid analysis.

We found that the *C. elegans* enzyme is active at the physiological growing temperatures for the nematode,³¹ while

the human enzyme is most active at the body’s expected physiological temperature of 37 °C.²⁶ These PRMT9 enzymes are active over a range of temperatures wider than that of the PRMT7 enzymes analyzed in Figure 3.

THW Loop Residues Are Responsible for the SDMA Forming Ability of *C. elegans* PRMT-9. The central residue in the conserved THW loop has been shown to play a role in the formation of MMA and SDMA in *T. brucei* PRMT7 and in human PRMT9.²³ Specifically, mutating the central cysteine residue in the human PRMT9 THW motif to histidine greatly enhanced the ability of the enzyme to form MMA and weakened its ability to form SDMA.²² Making the same mutation in the *C. elegans* PRMT-9 enzyme, converting the alanine residue in the central position of the THW loop to a histidine residue, resulted in the loss of SDMA formation and reduced the extent of MMA formation (Figure 12). The inverse H300A mutation made in the *C. elegans* PRMT-7 enzyme resulted in the loss of activity (data not shown). Interestingly, additional DTT added to the reaction mixtures, for the wild-type and A391H mutant enzymes (blue line in all panels), enhanced the formation of MMA but did not affect the formation of SDMA (Figure 12).

■ DISCUSSION

While many different physiological roles have been described for PRMT7,^{10–16,30} endogenous *in vivo* substrates have yet to be identified. PRMT7 is widely distributed across vertebrates, invertebrates, and plants. Fungi such as *Saccharomyces cerevisiae* and *Endocarpon pusillum* do not contain a PRMT7 ortholog, as their top BLAST hits match well with the major PRMT1 ortholog (Figure 1). Other studies have not found corresponding orthologs in *S. cerevisiae* or *Schizosaccharomyces pombe*.¹ What was most surprising is our identification of potential bacterial PRMT7 orthologs from *R. leguminosarum* and *G. sulfurreducens* that were mutual best hits to human PRMT7. Methylation of arginine residues has never been observed in bacteria, and no PRMT genes have been previously reported in bacteria. These prospective bacterial orthologs lack an apparent THW loop and do not contain the ancestrally duplicated C-terminal domain, much like the *T. brucei* ortholog of PRMT7. *G. sulfurreducens* does contain the characteristic TPR motifs of human PRMT9, yet the closest human hit when analyzed with protein BLAST is PRMT7. Further work needs to be done to confirm if these potential enzymes actually methylate arginine residues. Presumably, the retention of PRMT7 in higher organisms reflects its specific cellular roles, including the limitation of its activity to monomethylation, its substrate specificity, and its potential for regulation and crosstalk with other modifying enzymes.

Optimizing reaction conditions for the *C. elegans* PRMT-7 enzyme, such as lowering the reaction temperature, removing metal ion chelators, and adding reducing agents such as DTT, proved to be essential for determining its substrate preference and methylation activity. While the optimal temperature for the activity of *C. elegans* PRMT-7 falls clearly in the physiological range of this organism, the human PRMT7 enzyme also demonstrates a similar temperature preference range where it is much more active than it is at 37 °C. This is an unusual property for a human enzyme and may be important for its activity in regions of localized hypothermia such as in lung tissue exposed to cold air or in the extremities. It would be interesting to correlate the function of PRMT7 with the expression of cold-inducible RNA binding proteins CIRP and





Figure 11. Temperature dependence of *C. elegans* PRMT-9. Methylation reaction mixtures consisting of 5 μg of substrate (*H. sapiens* GST-tagged SF3B2 fragment 401–550) were reacted with 2 μg of enzyme (both *C. elegans* and *H. sapiens* PRMT9 enzymes) at the various temperatures indicated for 20 h in 50 mM potassium HEPES and 10 mM NaCl (pH 8.2) with 0.7 μM [methyl- ^3H]AdoMet in a final reaction volume of 60 μL . Reactions were quenched by the addition of sample loading buffer, and proteins were separated using SDS–PAGE using the same methods described in the legend of Figure 3. The gel was dried and exposed to autoradiography film for 21 days at -80°C . A Coomassie gel (top) and fluorograph (bottom) are shown. Molecular weight positions are shown from approximately 2 μg of unstained SDS–PAGE broad range marker as described in the legend of Figure 3. This experiment was replicated independently, and a third replicate using the *C. elegans* PRMT-9 enzyme was performed. The bottom panel indicates the densitometry of the bands comparing the methylation activity of the *C. elegans* enzyme to that of the human enzyme. Densitometry was performed using ImageJ on the scanned images of the films and quantified as described in the legend of Figure 3 (*C. elegans*, red symbols; human, blue symbols). A solid line is drawn for the average of the normalized values of the replicates. The optimal growth temperatures for *C. elegans*³¹ and humans are indicated.




Figure 12. *C. elegans* PRMT-9 THW loop residues are important for conferring SDMA specificity and a small DTT effect. (A) *C. elegans* PRMT-9 enzyme (2 μ g) wild type (top) or THW loop mutant A391H (bottom) was reacted with approximately 5 μ g of *C. elegans* SFTB-2 for 20 h at 25 °C in the reaction buffer containing 50 mM potassium HEPES and 10 mM NaCl (pH 8.0) with (blue line) or without (red line) 1 mM DTT, as indicated. Mutation of THW loop residue A391 to the conserved histidine residue abolishes the SDMA activity, producing only MMA, consistent with previous work with the human enzyme PRMT9 C431H mutant.²² These reactions were single replicates.

RBM3, both of which are known to be methylated at arginine residues.³³ Alternatively, the small amount of activity of human PRMT7 at 37 °C may be sufficient for its function, or there may be other proteins or small molecules present in human cells that allow for enhanced activity at normal body temperature.

We show here that *C. elegans* PRMT-7 methylates human histone H2B peptides, prime *in vitro* substrates for the human PRMT7 enzyme, but less methylation of the corresponding *C. elegans* H2B peptide was seen (Figure 4A). This result suggests that histone H2B may not be a major physiological substrate for *C. elegans* PRMT-7. It has been shown in proteomic studies that there is divergence in conservation of methylation sites across various organisms,³⁴ indicating the methylation sites can differ across different species and organisms.

Our results also indicate the importance of multiple arginine residues in an R-X-R motif for recognition by both *C. elegans* PRMT-7 and human PRMT7. However, the *C. elegans* enzyme does not appear to have a strong preference for adjacent basic residues and methylated substrates with adjacent glycine residues such as pSmD3. Peptides with single arginine residues were generally not substrates, but a histone H4 1–16 peptide containing a single arginine residue was weakly methylated by *C. elegans* PRMT-7. It appears that the length of the peptide, the positioning of arginine residues, and the nature of adjacent and distant residues play a role in substrate specificity.

Comparison of the activity of the human and *C. elegans* PRMT7s on intact human recombinant histones confirms the specificity of the human enzyme for H2B.¹⁷ However, we observe here that *C. elegans* PRMT-7 methylates all of the histone proteins, with the exception of only weakly methylating histone H4 (Figure 6). This broader substrate methylation leads us to believe that its physiological substrates, while not identified yet, could potentially be different from and more numerous than those of its mammalian ortholog.

To further probe the substrate specificity of these enzymes, we examined the active site architectures of *C. elegans*,²⁴ *T. brucei*,²⁰ and *M. musculus* PRMT7.²⁵ Surface modeling of their crystal structures reveals differences that may relate to their activities. As discussed in Results, the distances between the two terminal glutamates in the double E loop were closer together in the *C. elegans* and *T. brucei* structures than in the mouse structure. In addition, there are differences in residues, in particular a phenylalanine (*C. elegans* or *T. brucei*) or serine (mouse) residue that comes in contact with the second glutamate of the double E loop. These structural differences may explain why we see such distinct substrate specificity among the various orthologs of PRMT7, indicating that the enzyme may have evolved to have different functions and substrates among different organisms. Further work needs to be done to confirm the importance of these residues in conferring substrate specificity.

C. elegans also has a PRMT-9 ortholog that has been conserved from invertebrates to mammals with a potentially similar physiological function. We show here that the *C. elegans* PRMT-9 enzyme produces SDMA and MMA from the splicing factor SF3B2 ortholog in *C. elegans*, SFTB-2. Additionally, we found that the enzyme methylates the same R508 residue on SF3B2 as the human enzyme, suggesting that PRMT-9 may also play a regulatory role in nematode alternative RNA splicing. It will now be important to directly examine the effect of methylation of SFTB-2 on the regulation of splicing in *C. elegans*, a process that has been shown to be similar to human splicing mechanisms.³⁵

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.biochem.7b00283.

Peptide sequences used in the text (Table S1) (PDF)

Accession Codes

C. elegans PRMT-7 (UniProt entry Q9XW42), *C. elegans* PRMT-9 (UniProt entry O02325), human PRMT7 (UniProt entry Q9NVM4), human PRMT9 (UniProt entry Q6P2P2), *C. elegans* SFTB-2 (UniProt entry O16997), human SF3B2 (UniProt entry Q13435), and amino acids 1–145 of human fibrillarin (UniProt entry P22087).

■ AUTHOR INFORMATION

Corresponding Author

*Department of Chemistry and Biochemistry and Molecular Biology Institute, University of California, Los Angeles, 607 Charles E. Young Dr. E., Los Angeles, CA 90095-1569. Telephone: 310-825-8754. E-mail: clarke@mbi.ucla.edu.

ORCID

Andrea Hadjikyriacou: 0000-0002-3420-367X

Steven G. Clarke: 0000-0002-7303-6632

Author Contributions

A.H. and S.G.C. conceived the project and wrote the manuscript. A.H. performed the experiments.

Funding

This work was supported, in whole or in part, by National Institutes of Health Grants GM026020 (to S.G.C.) and GM007185, a Ruth L. Kirschstein National Research Service Award (to A.H.), funds from the Elizabeth and Thomas Plott Chair in Gerontology of the UCLA Longevity Center (to S.G.C.), funds from a UCLA Dissertation Year Fellowship (to A.H.), and a UCLA Faculty Research Award (to S.G.C.).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors thank Alexander Patananan (University of California, Los Angeles) for his advice on *C. elegans* biology and Akiyoshi Fukamizu (University of Tsukuba, Tsukuba, Japan) for providing the cDNA clones of *C. elegans* GST-PRMT-7 and GST-PRMT-9. The authors also thank Adam Frankel (University of British Columbia, Vancouver, BC) for the R1 peptide and Mark T. Bedford (M. D. Anderson Cancer Center, Houston, TX) for the GST-SF3B2 plasmid. The authors also thank members of the Clarke laboratory, in

particular Jonathan Lowenson, Kanishk Jain, and Rebecca Warmack, for useful comments on the manuscript.

■ ABBREVIATIONS

PRMT, protein arginine methyltransferase; MMA, ω - N^G -monomethylarginine; ADMA, ω - N^G, N^G -asymmetric dimethylarginine; SDMA, ω - N^G, N^G -symmetric dimethylarginine; AdoMet, S-adenosyl-L-methionine; [methyl-³H]AdoMet, S-adenosyl[methyl-³H]-L-methionine; GAR, glycine- and arginine-rich domain of human fibrillarin.

■ REFERENCES

- (1) Wang, Y., Wang, J., Chen, C., Chen, Y., and Li, C. (2015) A novel BLAST-Based Relative Distance (BBRD) method can effectively group members of protein arginine methyltransferases and suggest their evolutionary relationship. *Mol. Phylogenet. Evol.* 84, 101–111.
- (2) Wang, Y. C., and Li, C. (2012) Evolutionarily conserved protein arginine methyltransferases in non-mammalian animal systems. *FEBS J.* 279, 932–945.
- (3) Fuhrmann, J., Clancy, K. W., and Thompson, P. R. (2015) Chemical biology of protein arginine modifications in epigenetic regulation. *Chem. Rev.* 115, 5413–5461.
- (4) Bedford, M. T., and Clarke, S. G. (2009) Protein arginine methylation in mammals: Who, what, and why. *Mol. Cell* 33, 1–13.
- (5) Yang, Y., Hadjikyriacou, A., Xia, Z., Gayatri, S., Kim, D., Zurita-Lopez, C., Kelly, R., Guo, A., Li, W., Clarke, S. G., and Bedford, M. T. (2015) PRMT9 is a Type II methyltransferase that methylates the splicing factor SAPI45. *Nat. Commun.* 6, 6428.
- (6) Auclair, Y., and Richard, S. (2013) The role of arginine methylation in the DNA damage response. *DNA Repair* 12, 459–465.
- (7) Yang, Y., and Bedford, M. T. (2013) Protein arginine methyltransferases and cancer. *Nat. Rev. Cancer* 13, 37–50.
- (8) Baldwin, R. M., Morettin, A., and Côté, J. (2014) Role of PRMTs in cancer: Could minor isoforms be leaving a mark? *World Journal of Biological Chemistry* 5, 115–29.
- (9) Likhite, N., Jackson, C. A., Liang, M., Krzyzanowski, M. C., Lei, P., Wood, J. F., Birkaya, B., Michaels, K. L., Andreadis, S. T., Clark, S. D., Yu, M. C., and Ferkey, D. M. (2015) The protein arginine methyltransferase PRMT5 promotes D2-like dopamine receptor signaling. *Sci. Signaling* 8, ra115.
- (10) Blanc, R. S., Vogel, G., Chen, T., Crist, C., and Richard, S. (2016) PRMT7 preserves satellite cell regenerative capacity. *Cell Rep.* 14, 1528–1539.
- (11) Ying, Z., Mei, M., Zhang, P., Liu, C., He, H., Gao, F., and Bao, S. (2015) Histone arginine methylation by PRMT7 controls germinal center formation via regulating Bcl6 transcription. *J. Immunol.* 195, 1538–47.
- (12) Yao, R., Jiang, H., Ma, Y., Wang, L., Wang, L., Du, J., Hou, P., Gao, Y., Zhao, L., Wang, G., Zhang, Y., Liu, D. X., Huang, B., and Lu, J. (2014) PRMT7 induces epithelial-to-mesenchymal transition and promotes metastasis in breast cancer. *Cancer Res.* 74, 5656–5667.
- (13) Kernohan, K. D., McBride, A., Xi, Y., Martin, N., Schwartzenuber, J., Dymont, D. A., Majewski, J., Blaser, S., Care4Rare Canada Consortium, Boycott, K. M., and Chitayat, D. (2017) Loss of the arginine methyltransferase PRMT7 causes syndromic intellectual disability with microcephaly and brachydactyly. *Clin. Genet.* 91, 708–716.
- (14) Baldwin, R. M., Haghandish, N., Daneshmand, M., Amin, S., Paris, G., Falls, T. J., Bell, J. C., Islam, S., and Côté, J. (2015) Protein arginine methyltransferase 7 promotes breast cancer cell invasion through the induction of MMP9 expression. *Oncotarget* 6, 3013–3032.
- (15) Karkhanis, V., Wang, L., Tae, S., Hu, Y. J., Imbalzano, A. N., and Sif, S. (2012) Protein arginine methyltransferase 7 regulates cellular response to DNA damage by methylating promoter histones H2A and H4 of the polymerase δ catalytic subunit gene, POLD1. *J. Biol. Chem.* 287, 29801–29814.
- (16) Dhar, S. S., Lee, S. H., Kan, P. Y., Voigt, P., Ma, L., Shi, X., Reinberg, D., and Lee, M. G. (2012) Trans-tail regulation of MLL4

catalyzed H3K4 methylation by H4R3 symmetric dimethylation is mediated by a tandem PHD of MLL4. *Genes Dev.* 26, 2749–2762.

(17) Feng, Y., Maity, R., Whitelegge, J. P., Hadjikyriacou, A., Li, Z., Zurita-Lopez, C., Al-Hadid, Q., Clark, A. T., Bedford, M. T., Masson, J. Y., and Clarke, S. G. (2013) Mammalian protein arginine methyltransferase 7 (PRMT7) specifically targets RXR sites in lysine- and arginine-rich regions. *J. Biol. Chem.* 288, 37010–37025.

(18) Feng, Y., Hadjikyriacou, A., and Clarke, S. G. (2014) Substrate specificity of human protein arginine methyltransferase 7 (PRMT7) - The importance of acidic residues in the double E loop. *J. Biol. Chem.* 289, 32604–32616.

(19) Fisk, J. C., Sayegh, J., Zurita-Lopez, C., Menon, S., Presnyak, V., Clarke, S. G., and Read, L. K. (2009) A type III protein arginine methyltransferase from the protozoan parasite *Trypanosoma brucei*. *J. Biol. Chem.* 284, 11590–11600.

(20) Wang, C., Zhu, Y., Caceres, T. B., Liu, L., Peng, J., Wang, J., Chen, J., Chen, X., Zhang, Z., Zuo, X., Gong, Q., Teng, M., Hevel, J. M., Wu, J., and Shi, Y. (2014) Structural determinants for the strict monomethylation activity by *Trypanosoma brucei* protein arginine methyltransferase 7. *Structure* 22, 756–768.

(21) Debler, E. W., Jain, K., Warmack, R. A., Feng, Y., Clarke, S. G., Blobel, G., and Stavropoulos, P. (2016) A glutamate/aspartate switch controls product specificity in a protein arginine methyltransferase. *Proc. Natl. Acad. Sci. U. S. A.* 113, 2068–2073.

(22) Jain, K., Warmack, R. A., Debler, E. W., Hadjikyriacou, A., Stavropoulos, P., and Clarke, S. G. (2016) Protein arginine methyltransferase product specificity is mediated by distinct active-site architectures. *J. Biol. Chem.* 291, 18299–18308.

(23) Takahashi, Y., Daitoku, H., Yokoyama, A., Nakayama, K., Kim, J.-D., and Fukamizu, A. (2011) The *C. elegans* PRMT-3 possesses a type III protein arginine methyltransferase activity. *J. Recept. Signal Transduction Res.* 31, 168–172.

(24) Hasegawa, M., Toma-Fukai, S., Kim, J. D., Fukamizu, A., and Shimizu, T. (2014) Protein arginine methyltransferase 7 has a novel homodimer-like structure formed by tandem repeats. *FEBS Lett.* 588, 1942–1948.

(25) Cura, V., Troffer-Charlier, N., Wurtz, J.-M., Bonnefond, L., and Cavarelli, J. (2014) Structural insight into arginine methylation by the mouse protein arginine methyltransferase 7: a zinc finger freezes the mimic of the dimeric state into a single active site. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 70, 2401–2412.

(26) Hadjikyriacou, A., Yang, Y., Espejo, A., Bedford, M. T., and Clarke, S. G. (2015) Unique features of human protein arginine methyltransferase 9 (PRMT9) and its substrate RNA splicing factor SF3B2. *J. Biol. Chem.* 290, 16723–16743.

(27) Zurita-Lopez, C. I., Sandberg, T., Kelly, R., and Clarke, S. G. (2012) Human protein arginine methyltransferase 7 (PRMT7) is a type III enzyme forming ω -N^G-monomethylated arginine residues. *J. Biol. Chem.* 287, 7859–7870.

(28) Tang, J., Gary, J. D., Clarke, S., and Herschman, H. R. (1998) PRMT 3, a type I protein arginine N-methyltransferase that differs from PRMT1 in its oligomerization, subcellular localization, substrate specificity, and regulation. *J. Biol. Chem.* 273, 16935–16945.

(29) McWilliam, H., Li, W., Uludag, M., Squizzato, S., Park, Y. M., Buso, N., Cowley, A. P., and Lopez, R. (2013) Analysis tool web services from the EMBL-EBI. *Nucleic Acids Res.* 41, W597–W600.

(30) Jelinic, P., Stehle, J. C., and Shaw, P. (2006) The testis-specific factor CTCFL cooperates with the protein methyltransferase PRMT7 in H19 imprinting control region methylation. *PLoS Biol.* 4, e355.

(31) Corsi, A. K., Wightman, B., and Chalfie, M. (2015) A transparent window into biology: A primer on *Caenorhabditis elegans*. In *Worm Book Genetics* (De Stasio, E. A., Ed.) pp 387–407, Worm Book.

(32) Petrella, L. N. (2014) Natural variants of *C. elegans* demonstrate defects in both sperm function and oogenesis at elevated temperatures. *PLoS One* 9, e112377.

(33) LLeonart, M. E. (2010) A new generation of proto-oncogenes: Cold-inducible RNA binding proteins. *Biochim. Biophys. Acta, Rev. Cancer* 1805, 43–52.

(34) Larsen, S. C., Sylvestersen, K. B., Mund, A., Lyon, D., Mullari, M., Madsen, M. V., Daniel, J. A., Jensen, L. J., and Nielsen, M. L. (2016) Proteome-wide analysis of arginine monomethylation reveals widespread occurrence in human cells. *Sci. Signaling* 9, rs9.

(35) Gracida, X., Norris, A. D., and Calarco, J. A. (2016) Regulation of tissue-specific alternative splicing: *C. elegans* as a model system. In *Advances in Experimental Medicine and Biology*, pp 229–261, Springer International Publishing Switzerland, Cambridge, MA.

(36) Gottschling, H., and Freese, E. (1962) Tritium isotope effect on ion exchange chromatography. *Nature* 196, 829–831.